

Retinitis pigmentosa is a group of genetic disorders that affect the retina's ability to respond to light.

The retina is the layer of light-sensing cells lining the back of your eye that converts light rays into impulses. The impulses are sent through the optic nerve to your brain, where they are recognized as images.

WHAT ARE THE SYMPTOMS OF RETINITIS PIGMENTOSA?


Retinitis pigmentosa causes slow loss of vision. Symptoms begin with decreased night vision and later progress to loss of peripheral (side) vision. Some people may also have difficulty identifying colors. The rate of vision change varies in different people depending on the genetic makeup of their disorder.

As night vision decreases, the ability to adjust to darkness becomes slower and slower. You may:


- stumble over objects in the dark;
- find driving at dusk and night difficult;
- see poorly in dimly lit rooms, such as a movie theater.

While your vision during the day may be completely normal, your inability to see in dark conditions is considered "night blindness."

Loss of side vision also makes moving around more difficult, causing noticeable clumsiness from not being able to see objects below and around you. As the outer areas of vision slowly disappear, the condition commonly referred to as "tunnel vision" occurs. Many people retain a wide enough scope of useful vision for a long period of time. In some cases, central vision



Normal retina



Retina with retinitis pigmentosa

may be affected first, making detail work difficult, such as reading or threading a needle. This is referred to as **macular dystrophy**, because the central area of the retina, called the **macula**, is affected.

WHAT CAUSES RETINITIS PIGMENTOSA?

Retinitis pigmentosa is thought to be caused by defects of certain genes in the body. The defects cause cells in the retina to malfunction and begin to break down.



retinitis pigmentosa

Retinitis pigmentosa is often hereditary (runs in families). If you or your partner have retinitis pigmentosa, there may be up to a 50 percent chance that you will pass it along to your children. Ask your ophthalmologist (Eye M.D.) about genetic counseling if you are planning to have children.

AMERICAN FOUNDATION FOR THE BLIND

11 Penn Plaza, Suite 300 New York, NY 10001 800.232.5463 www.afb.org

IS THERE ANY TREATMENT FOR RETINITIS PIGMENTOSA?

Research has shown that Vitamin A can slow the progression of retinitis pigmentosa. Your ophthalmologist can advise you about the risks and benefits of Vitamin A and how much you can take safely.

Research is currently directed toward finding the hereditary cause of many types of retinitis pigmentosa. As hereditary defects are discovered, it may be possible to develop treatments to prevent the progression of retinitis pigmentosa.

Special low-vision aids and services can help people with poor vision continue doing many daily-life activities. Ask your ophthalmologist or contact the following organizations for more information and resources near you:

NATIONAL ASSOCIATION FOR VISUALLY HANDICAPPED

NAVH New York City 22 West 21st Street, 6th Floor New York, NY 10010 212.889.3141 www.navh.org

NAVH San Francisco 507 Polk Street, Suite 420 San Francisco, CA 94102 415.775.6284 www.navh.org

Academy reviewed 10/09

COMPLIMENTS OF YOUR OPHTHALMOLOGIST:

057169

© 2009 American Academy of Ophthalmology. The American Academy of Ophthalmology, The Eye M.D. Association and the Academy logo are registered trademarks of the American Academy of Ophthalmology.





ISBN 978-1-61525-052-3